

PERACETIC ACID AND SODIUM HYPOCHLORITE EFFECTIVENESS IN REDUCING RESISTANT STAGES OF MICROORGANISMS

Briancesco R., Veschetti E., Ottaviani M., Bonadonna L.

Department Environment and Primary Prevention, Istituto Superiore di Sanità, Roma, Italy

SUMMARY

A comparative study on the efficacy of peracetic acid and sodium hypochlorite in inactivating resistant stages of microorganisms such as *Cryptosporidium*, *Giardia* and *Cl. perfringens* was carried out. Furthermore the evaluation of the potential reciprocal correlation among the concentrations of the organisms was performed. The results obtained indicate that, at the same experimental conditions, peracetic acid and sodium hypochlorite have nearly similar reduction power against the resistant stages of *Giardia* and *Cl. perfringens*. Both the oxidants are instead less efficient in the abatement of *Cryptosporidium* oocysts. Findings have also confirmed our previous studies on the absence of association between *Clostridium* and the protozoa.

Key words: *Clostridium perfringens*, protozoa, peracetic acid, sodium hypochlorite, wastewater

Address for correspondence: L. Bonadonna, Istituto Superiore di Sanità, Viale Regina Elena, 299, 00161 Roma, Italy. E-mail: lucybond@iss.it

INTRODUCTION

Pathogens in water are generally considered a higher health risk than chemicals (1). Current drinking water disinfection practices provide the means to control most pathogenic microorganisms responsible for the major waterborne diseases. Most outbreaks in recent years have been caused by viruses and protozoa that are generally more resistant to chlorine-based disinfection than are pathogenic bacteria, the primary targets of concern in the past decades. Given the refractory nature of the protozoa *Giardia* and *Cryptosporidium* to standard water chlorination, significant effort has been expended to find new chemical agents, combinations, or conditions that could be used to effectively treat water. Furthermore, chlorine is also known to raise serious toxic effects on living organisms and several studies have demonstrated toxic, mutagenic and carcinogenic properties of its by-products (2, 3).

In the last years, peracetic acid has been introduced as disinfectant in agronomical and alimentary industries as well as in health sector, and only recently for wastewater disinfection (4, 5, 6). Its major advantages are the ease of implementing treatment, broad spectrum of activity even in the presence of heterogeneous organic matter and absence of persistent toxic or mutagenic residuals or by-products (7, 8).

Till now no data on the effects of peracetic acid on resistant stages, i.e. cysts and/or spores, of microorganisms in wastewater are available.

Taking in mind these issues, the current study was aimed at following: i) to monitor concentrations of the infective and resistant stages, oocysts and cysts, respectively, of *Cryptosporidium* and *Giardia*, in wastewater after a coarse screening and in the effluent after a secondary clarification in a municipal wastewater-treatment plant, ii) to estimate the efficacy of these treatments in removing *Cryptosporidium* oocysts, *Giardia* cysts and *Clostridium perfringens* spores, iii) to compare simultaneously the efficacy

of two disinfectants (peracetic acid and sodium hypochlorite) in inactivating *Cryptosporidium* oocysts, *Giardia* cysts and *Cl. perfringens* spores during tertiary treatment, iv) to evaluate the potential reciprocal correlation among the concentrations of enteric protozoa oo/cysts and *Cl. perfringens* spores as putative indicator of their presence.

MATERIALS AND METHODS

Processing of sewage samples. Wastewater samples were obtained from a municipal plant located in Rome (Italy). The experimentation was partially carried out in a pilot plant installed in the municipal plant that uses a conventional wastewater-treatment system consisting of screening, primary clarification, aeration and biological oxidation through activated sludge, secondary clarification, and chlorination. The study was carried out on raw water (only subjected to a coarse screening), secondary effluent and two different tertiary effluents coming from alternative disinfection processes at the pilot plant.

Description of the pilot plant. The pilot plant consisted of a metallic structure subdivided in two parallel tanks. Each tank was employed for the contact between sewage and a disinfectant. Both the tanks were fed by the effluent from the secondary settler placed after the activated-sludge biological tank of the municipal plant. The effluent sucked by a submersible pump was introduced in the tanks by means of a pipe divided by a tee joint and a magnetic volume meter and a throttle valve were installed along each breach to obtain an independent control of the flow in the two tanks.

Residence times (t_r) of each tank were determined by injecting a tracer (6 l of a saturated solution of sodium chloride) into the sewage stream at the inlet of the pilot plant and by measuring the conductivity at the exit of the tanks. These measures were carried out at two different flow rates (2.0 and 8.0 l/s). The average ratio

Table 1. Concentration¹⁾

Sample N°	Water after coarse screening			Water after secondary clarification			Reduction (%)		
	Cysts/l	Oocysts/l	Spores/l	Cysts/l	Oocysts/l	Spores/l	Cysts	Oocysts	Spores
1	1.5×10 ⁴	1.2×10 ³	1.4×10 ⁶	1.5×10 ³	5.0×10 ²	6.0×10 ⁴	90	58	96
2	3.1×10 ³	< 1.0×10 ⁰	6.0×10 ⁵	1.7×10 ³	< 1.0×10 ⁰	3.1×10 ⁴	45	–	95
3	3.0×10 ⁴	6.0×10 ²	1.4×10 ⁵	4.3×10 ³	1.5×10 ²	2.7×10 ⁴	86	75	81
4	5.7×10 ³	4.5×10 ²	3.0×10 ⁵	6.2×10 ²	2.1×10 ²	7.3×10 ⁴	93	53	76
5	2.7×10 ³	3.0×10 ³	n.d.	1.4×10 ³	1.3×10 ³	n.d.	48	57	–
6	1.2×10 ⁴	1.0×10 ³	2.5×10 ⁵	3.0×10 ²	3.9×10 ²	1.3×10 ⁴	98	61	95
7	2.5×10 ⁴	1.0×10 ³	2.0×10 ⁴	3.1×10 ³	4.3×10 ²	1.4×10 ³	88	57	93
8	3.5×10 ⁴	8.0×10 ²	5.0×10 ⁵	2.1×10 ³	2.2×10 ²	8.0×10 ³	94	64	98
9	2.5×10 ⁴	< 1.0×10 ⁰	1.6×10 ⁶	2.5×10 ³	< 1.0×10 ⁰	n.d.	90	–	–

n.d.: not determined

between t_R and the time necessary to fill each tank was 0.78 for both contact chambers. The value of the actual contact time at varying of the flow rate was calculated by multiplying the average ratio by geometric volume of the fluid and the volumetric flow rate measured during each disinfection test.

Disinfection. Sodium hypochlorite (HYP) and peracetic acid (PAA) were tested in parallel during the tertiary treatment of the secondary effluent. The experiments were performed in the following conditions: disinfectant concentrations ranging from 2.0 to 4.5 mg/l, contact time of 20–30 min, pH 7 and temperature of 20 °C. Disinfectants were introduced in the respective tanks by two laboratory-made pneumatic pumps. The concentration of the disinfectants was determined daily before starting tests. Iodometric procedures were employed to measure levels of free chlorine and peroxiacid in sodium-hypochlorite and peracetic acid solutions, respectively. Free iodine liberated from KI by HYP or PAA was titrated against standardised Na₂S₂O₃ solution (0.1N). The interference caused by H₂O₂ in the determination of PAA titre was eliminated by adding some drops of catalase. Other analytical procedures were performed according to the methods described in the literature (9, 10).

Cryptosporidium and Giardia oocysts detection. The detection was carried out on adequate volumes of the following samples:

- raw water collected after a coarse screening;
- secondary effluent (water coming from secondary clarification, before entering the pilot plant);
- water running off the pilot plant after the treatment with sodium hypochlorite;
- water running off the pilot plant after the treatment with peracetic acid.

A total of 32 samples were collected in plastic tanks. Sodium thiosulfate (10 mg/l) and catalase (0.1 mg/l) were added to disinfected samples to inactivate any residual biocide. Cellulose-acetate membrane filters (Millipore, USA) were utilized to entrap *Cryptosporidium* oocysts and *Giardia* cysts from water samples and processed following the membrane filter dissolution method, according to Aldom and Chagla (11) and Graczyk and coll. (12). Sheather's sucrose density centrifugation was used to

separate cysts and oocysts from suspended matter in the pellet (13). Parasites were detected by direct immunofluorescence with anti-*Giardia* and anti-*Cryptosporidium* monoclonal antibodies (Meridian Diagnostics, USA) taking into consideration morphology, size and color of the particles with respect to a positive standard. The presumed inactivation of cysts and oocysts was evaluated through the inspection of internal structures by contrast phase microscopy.

Clostridium perfringens spores detection. Samples were pasteurized by heating to 75 °C for 15 min; spores were enumerated on OPSP (Perfringens Agar Base, Oxoid, UK) after incubation under anaerobic condition for 48 h. Black colonies were confirmed by the miniaturised system API 20A (Biomérieux, France).

Statistical analysis. Results were analysed with the statistical package SPSS v. 10.0.

RESULTS

In Table 1 the concentrations of *Giardia* cysts, *Cryptosporidium* oocysts and *Cl. perfringens* spores in water after the coarse screening and in the effluent after the secondary clarification are reported. The percentage of reduction was also calculated. As it was expected higher values were observed for *Cl. perfringens* spores, followed by *Giardia* cysts and *Cryptosporidium* oocysts that, contemporaneously, showed the highest variability. Their mean reduction after the clarification treatment was 81%, 61% and 91%, respectively. The extent of reduction was statistically significant for *Giardia* cysts ($p < 0.01$) and *Cl. perfringens* spores ($p < 0.01$).

In both the kind of water, no significative reciprocal correlation was observed among the three parameters.

Table 2 shows the reduction in the target organisms caused by the two alternative disinfection procedures. At the same c:t values, expressed as mg·min/l, sodium hypochlorite and peracetic acid exhibited nearly similar trends, with more uniform reduction values of oocysts and spores. The t-test, calculated with respect to the microorganism densities at c:t of 120 mg·min/l, has demonstrated that the concentration of *Giardia* cysts and *Cryptosporidium*

Table 2. Reduction of *Giardia* cysts, *Cryptosporidium* oocysts and *Cl. perfringens* spores after the two different disinfection processes at the pilot plant

Sample N°	Water treated with sodium hypochlorite			Water treated with peracetic acid		
	2	2	3	2	2	3
C . t (mg . min/l)	60	90	120	60	90	120
Reduction (%) of <i>Giardia</i> cysts	94.0	97.0	98.0	92.8	96.5	91.6
Reduction (%) of <i>Cryptosporidium</i> oocysts	73.0	74.0	68.4	71.0	75.0	74.5
Reduction (%) of <i>Cl. perfringens</i> spores	99.3	99.8	n.d.	97.4	99.7	n.d.

n.d.: not determined

Table 3. Survival of *Giardia* cysts, *Cryptosporidium* oocysts and *Cl. perfringens* spores in the post-disinfected effluents

	Mean percentage of surviving organisms*					
	Sodium hypochlorite treatment			Peracetic acid treatment		
C . t (mg . min/l)	60	90	120	60	90	120
N°**	2	2	3	2	2	3
<i>Giardia</i> cysts	49	43	48	57	50	78
<i>Cryptosporidium</i> oocysts	75	73	77	80	76	71
<i>Cl. perfringens</i> spores	3	7.5	n.d.	28	31	n.d.

*Post-disinfected /pre-disinfected densities ($\times 100$); **N°: number of paired pre- and post-disinfected samples

n.d.: not determined

oocysts in the effluents treated with sodium hypochlorite was not significantly different from that present in the effluent treated with peracetic acid.

Values of the survival mean percentage of protozoan oo/cysts and *Cl. perfringens* spores in the post-disinfected effluent in respect to their densities in the pre-disinfected effluent are reported in Table 3. Different survival values were calculated for *Cl. perfringens* with respect to *Giardia* and *Cryptosporidium*. In fact, spores appeared to be more affected by both the disinfectants than *Giardia* and *Cryptosporidium* and much more by sodium hypochlorite than peracetic acid. The total mean percentage of spore inactivation was 95% and 71%, respectively, when sodium hypochlorite and peracetic acid were used. For *Giardia* cysts and *Cryptosporidium* oocysts, less influenced by both the disinfectants, the calculated values reached the 53% and 38% and the 25% and 24%, respectively.

DISCUSSION AND CONCLUSIONS

Data collected during this investigation indicate that high numbers of oocysts, cysts and spores were detected in water treated at the municipal wastewater-treatment plant. All samples were positive for cysts and spores, whereas the 22% of the samples gave negative results for oocysts. These findings confirm other studies

conducted in Italy where *Giardia* is detected more frequently in wastewater and at higher concentrations than *Cryptosporidium* (14). The lower *Cryptosporidium* density in waste water could be due to its probable actual lower occurrence in water in general but also due to methodological difficulties in oocysts detection caused by its small size. In fact, taking into consideration the recovery efficiency of the method applied to raw water that, for *Cryptosporidium* oocysts, corresponds to the 25–40%, it is presumable an underestimation of its concentrations.

Variable values of reduction of the resistant stages of both the protozoa were calculated according to the concentrations obtained after the secondary clarification. Values ranging from the 45% to 98% for *Giardia* and from the 53% to 75% for *Cryptosporidium* were recorded. They could be attributed to the changeable turbidity values measured in the analyzed waters ($4\div 18$ NTU). In fact, higher water turbidity values can interfere with the protozoa recovery (15). Besides, while the extent of reduction was statistically significative for *Giardia* and *Cl. perfringens*, it was not for *Cryptosporidium*. Once more, it could be ascribed to the larger size of cysts and spores, probably more affected by the sedimentation forces to which the effluent was subjected during the secondary clarification treatment.

The results obtained at the pilot plant indicate that, under the same experimental conditions, peracetic acid and sodium hypochlorite have nearly similar reduction power against the resistant stages of *Giardia* and *Cl. perfringens*. Both the oxidants are less efficient, and equivalent, in the abatement of *Cryptosporidium* oocysts, also at the highest dose (concentration 4.5 mg/l, contact time 30 min), thus confirming their greater resistance to biocides.

As far as the possible role of *Cl. perfringens* spores as indicator of occurrence of protozoan oo/cysts in treated water is concerned, our results evidenced the absence of association between *Clostridium* and the protozoa. This lack of consistent relationship in the examined water strengthen once again data previously find by the authors (16): the numbers of *Cl. perfringens* cannot be used to assess the risk of *Cryptosporidium* and *Giardia*.

Investigations of commonly used water treatment technologies indicate that both the protozoa are highly resistant to water treatment processes (17). Physical methods (microfiltration and UV irradiation) or sequential treatment using different disinfection agents have been found to be useful in certain circumstances (18). For this reason, in relation to the preliminary results obtained in this investigation, further studies are planned to establish the effective doses of peracetic acid necessary to give satisfactory reduction in protozoan numbers in sewage effluents under operational conditions and to identify its potential by-products.

REFERENCES

1. Craun GF: Safety of water disinfection: balancing chemical and microbial risks. Washington (USA): International Life Science Institute Press; 1993.
2. Glaze WH, Henderson JE: Formation of organochlorine compounds from the chlorination of a municipal secondary effluents. J Wat Pollut Control Fed. 1975;47:2511–5.
3. Rudd T, Hopkinson LM: Comparison of disinfection techniques for sewage and sewage effluents. J Instit Wat Environ Man. 1989;3: 612–8.
4. Lefevre F, Audic JM, Ferrand F: Peracetic acid disinfection of secondary effluents discharged off coastal seawater. Wat Sci Tech. 1992;25:155–64.

5. Baldry MGC, French MS, Slader D: The activity of peracetic acid on sewage indicator bacteria and viruses. *Wat Sci Tech.* 1991;24(2):353–7.
6. Booth RA, Lester JN: The potential formation of halogenated by-products during peracetic acid treatment of final sewage effluent. *Water Res.* 1995;29(7):1793–801.
7. Morris R: Reduction of microbial levels in sewage effluents using chlorine and peracetic acid disinfectants. *Water Sci Tech.* 1993;27(3-4):387–93.
8. Veschetti E, Cutilli D, Bonadonna L, Della Libera S, Ottaviani M, Martini C, Alaimo C, Pizzari P, Anastasi P: Preliminary results on the possibility of using peracetic acid as disinfectant of wastewater. In: *Advanced Wastewater Treatment, Recycling and Reuse: Proceedings of 2nd International Conference*; 1998 Sep 14–16; Milano, Italy. p. 397–404.
9. *Metodi Analitici per le acque*. Roma: Istituto di Ricerca sulle Acque, Consiglio Nazionale delle Ricerche; 1994.
10. American Public Health Association; Water Environmental Federation: *Standard Methods for the Examination of Water and Wastewater*. 19th ed. Washington DC: American Water Works Association; 1995.
11. Aldom JE, Chagla AH: Recovery of *Cryptosporidium* oocysts from water by a membrane filter dissolution method. *Appl Environ Microbiol.* 1995;20:186–7.
12. Graczyk TK, Cranfield MR, Fayer R: Recovery of waterborne oocysts of *Cryptosporidium* from water samples by membrane-filter dissolution method. *Paras Res.* 1997;83:121–5.
13. Briancesco R, Bonadonna L: Metodi per il rilevamento di cisti e oocisti di protozoi patogeni nelle acque superficiali. In: *Notiziario dei Metodi Analitici*. Roma: Istituto di ricerca sulle acque, CNR; 2000. p. 20–6.
14. Bonadonna L, Briancesco R, Cataldo C, Divizia M, Donia D, Panà A: Fate of bacterial indicators, viruses and protozoan parasites during a wastewater multi-component treatment system. *New Microbiol.* 2002; 25:413–20.
15. Fricker CR, Medema GD, Smith HV: Protozoan parasites (*Cryptosporidium*, *Giardia*, *Cyclospora*). In: *Microbial agents in drinking water: addendum to the 2nd edition of the Guidelines for drinking water quality*. Geneva: World Health Organization; 1999.
16. Briancesco R, Bonadonna L: An Italian study on *Cryptosporidium* and *Giardia* in wastewater, fresh water and treated water. *Environ Monit Assess.* 2005;104:445–57.
17. Robertson LJ, Smith HV, Paton CA: Occurrence of *Giardia* cysts and *Cryptosporidium* oocysts in sewage influent in six treatment plants in Scotland and prevalence of cryptosporidiosis and giardiasis diagnosed in the communities served by plants. In: Betts WB, Casemore DP, Fricker CR, Smith HV, Watkins J, editors. *Protozoan parasites and Water*. Cambridge: Royal Society of Chemistry; 1995. p. 47–9.
18. Fricker CR, Crabb JH: Water-borne cryptosporidiosis: detection methods and treatment options. *Adv Parasitol.* 1998;40:241–78.

Revised and accepted February 22, 2005